



京天成生物技术（北京）有限公司

AbMax Biotechnology Co., LTD

北京海淀区上地东路 29 号 2B 702

Building-2B, Suite 702, 29 E ShangDi Road,

Beijing, 100085, China

Tel: 86-010-62980517, Fax: 86-010-62977292

E-mail: info@antibodychina.com

Anti-GST, Clone 4G4

(mouse monoclonal IgG)

Catalog # 05-0017

Immunogen: GST recombinant protein

Specificity: Recognizes GST tag.

Species Cross-reactivity: Not determined.

Formulation: 0.1mg of mouse monoclonal IgG in 0.1ml of 0.01M PB, pH 7.4, 25% glycerol, 0.02% sodium azide. Protein was determined by OD₂₈₀ absorbance.

Physical State: Frozen liquid


Storage: Store the product at -20°C. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid. Prepare working dilution fresh each day. Avoid repeated freezing and thawing. For maximum recovery of product, centrifuge the original vial prior to removing the cap.

**FOR IN VITRO RESEARCH USE ONLY
NOT FOR USE IN HUMANS**

Research Applications

ELISA: A working concentration of 0.5~2µg/ml is suggested. Detection sensitivity 0.8ng/ml.

Immunoblot Analysis: 0.5~2µg/ml.



GST recombinant protein was resolved by electrophoresis, transferred to PVDF membrane and reacted with the recommended dilution of Anti-GST. The membrane was incubated using a goat anti-mouse secondary antibody conjugated to HRP and in ECL detection system to visualize results. Arrow indicates different GST recombinant proteins.

Immunoblot Protocol

1. Perform SDS-polyacrylamide gel electrophoresis (SDS-PAGE) on samples and transfer the proteins to a PVDF membrane.
2. Block the blotted PVDF membrane in freshly prepared PBS containing 5% nonfat dry milk and 0.02% Tween-20 for 1 hour at room temperature or overnight at 4°C in a shaker.
3. Incubate the PVDF with the recommended dilution of Anti-GST diluted in freshly prepared 1% nonfat dry milk and 0.02% Tween-20 for 2 hour at room temperature in a shaker.
4. Wash the PVDF 3 times with PBST.
5. Incubate the PVDF in the secondary reagent of choice (a goat anti-mouse secondary antibody conjugated to HRP, Catalog # 02-0001, 1:4000 dilution was used) in 1% nonfat dry milk and 0.02% Tween-20 for 1 hour at room temperature in a shaker.
6. Wash the PVDF 3-5 times with PBST.
7. Incubate the PVDF in ECL substrate for 1 min and cover the membrane by PE film.
8. In the darkroom, expose the PVDF to Autoradiography film for 5 s~ 5 min to visualize results.